☐ Fast Pass

lue Positive Sepsis not picked up by Sepsis Screening Tool

1 of the following Triggers:		Examples of Poor Perfusion
Hypothermic (< 96.8F) *Immediate sepsis huddle		Capillary Refill > 3 seconds
Hyperthermic (>100.4)		Skin temperature: Cool or Cold
Elevated WBC (if known)		Change in Mental Status
Bandemia (if known)		Lethargic
	<u> </u>	Confused
AND		Agitated
AND		Irritable
3 of the Following Triggers		Inconsolable
High Risk Diagnosis (see examples)		Hypotension
Tachycardia		
· · · · · · · · · · · · · · · · · · ·		Examples of High Risk
Poor Perfusion (see examples)		Sickle Cell
Tachypnea or Grunting		Oncology
		Implanted port or device
Sepsis Huddle Request Time:		Medically complex
RN Name:		Neonate < 28 days (>100.4 *Immediate Sepsis)
MD Name:		Transplant
Notes:		Asplenic
		Short Gut
Antibiotic start time:		Immunosuppressed
<ul><li>☐ Negative Screen</li><li>☐ Fast Pass</li><li>☐ Positive Sepsis not picked up by Sepsis Scre</li></ul>	ening Tool	Patient Sticker
☐ Fast Pass	ening Tool	Patient Sticker
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers:	ening Tool	Examples of Poor Perfusion
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers:	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4)	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4)  Elevated WBC (if known)	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused
Fast Pass Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4)  Elevated WBC (if known)	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated
Fast Pass Positive Sepsis not picked up by Sepsis Scre  PEPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known) Bandemia (if known)	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated Irritable
Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known) Bandemia (if known)  AND	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable
Fast Pass Positive Sepsis not picked up by Sepsis Scre  PEPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known)  Bandemia (if known)  AND  3 of the Following Triggers	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated Irritable
Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known) Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples)	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated Irritable Inconsolable Hypotension
Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known) Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable  Hypotension  Examples of High Risk
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known)  Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples)	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated Irritable Inconsolable Hypotension  Examples of High Risk Sickle Cell
Positive Sepsis not picked up by Sepsis Scre  EPSIS SCREENING TOOL  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known) Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable  Hypotension  Examples of High Risk  Sickle Cell  Oncology
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known)  Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples) Tachypnea or Grunting	ening Tool	Examples of Poor Perfusion Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated Irritable Inconsolable Hypotension  Examples of High Risk Sickle Cell Oncology Implanted port or device
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known)  Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples) Tachypnea or Grunting  Sepsis Huddle Request Time:	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable  Hypotension  Examples of High Risk  Sickle Cell  Oncology  Implanted port or device  Medically complex
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known)  Bandemia (if known)  AND  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples) Tachypnea or Grunting  Sepsis Huddle Request Time: RN Name:	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable  Hypotension  Examples of High Risk  Sickle Cell  Oncology  Implanted port or device  Medically complex  Neonate < 28 days (>100.4 *Immediate Sepsis)
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4)  Elevated WBC (if known)  Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples) Tachypnea or Grunting  Sepsis Huddle Request Time: RN Name: MD Name:	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable  Hypotension  Examples of High Risk  Sickle Cell  Oncology  Implanted port or device  Medically complex  Neonate < 28 days (>100.4 *Immediate Sepsis)  Transplant
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4) Elevated WBC (if known)  Bandemia (if known)  AND  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples) Tachypnea or Grunting  Sepsis Huddle Request Time: RN Name:	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds Skin temperature: Cool or Cold Change in Mental Status Lethargic Confused Agitated Irritable Inconsolable Hypotension  Examples of High Risk Sickle Cell Oncology Implanted port or device Medically complex Neonate < 28 days (>100.4 *Immediate Sepsis) Transplant Asplenic
Positive Sepsis not picked up by Sepsis Scre  **EPSIS SCREENING TOOL**  1 of the following Triggers: Hypothermic (< 96.8F) *Immediate sepsis huddle Hyperthermic (>100.4)  Elevated WBC (if known)  Bandemia (if known)  AND  3 of the Following Triggers High Risk Diagnosis (see examples) Tachycardia Poor Perfusion (see examples) Tachypnea or Grunting  Sepsis Huddle Request Time: RN Name: MD Name:	ening Tool	Examples of Poor Perfusion  Capillary Refill > 3 seconds  Skin temperature: Cool or Cold  Change in Mental Status  Lethargic  Confused  Agitated  Irritable  Inconsolable  Hypotension  Examples of High Risk  Sickle Cell  Oncology  Implanted port or device  Medically complex  Neonate < 28 days (>100.4 *Immediate Sepsis)  Transplant

## **SEPSIS HUDDLE TOOL**

Next Steps:

- MD and RN go to bedside for assessment
- MD determines Sepsis Huddle Outcome:

Outcome: (check one helow)



	outcomer (check one below)
	Suspected Infection: Low suspicion for invasive infection
	Suspected Sepsis: Without organ dysfunction
	Critical Sepsis: With organ or cardiovascular dysfunction

- MD enters orders into EPIC Order set # 11408: PEDS ED SEPSIS
- MD to ask ESR to Everbridge page:
  - o CODE SEPSIS Pharmacist only to prepare antibiotic therapy.
  - o PPR if pharmacist is needed at bedside for vasopressors and RT is needed for intubation

Each provider is responsible for reviewing 100% of the Sepsis Screening Tool for their patients.  This review is to make sure that the initial decisions were appropriate to the final patient outcomes.					
MD Initials:	Time Zero:	Date:			

88-4600-1 (Rev. 4/21)

## **SEPSIS HUDDLE TOOL**

Next Steps:

- MD and RN go to bedside for assessment
- MD determines Sepsis Huddle Outcome:



X	Outcome: (check one below)
	Suspected Infection: Low suspicion for invasive infection
	Suspected Sepsis: Without organ dysfunction
	Critical Sepsis: With organ or cardiovascular dysfunction

- MD enters orders into EPIC Order set # 11408: PEDS ED SEPSIS
- MD to ask ESR to Everbridge page:
  - o CODE SEPSIS Pharmacist only to prepare antibiotic therapy.
  - o PPR if pharmacist is needed at bedside for vasopressors <u>and</u> RT is needed for intubation

·	Each provider is responsible for reviewing 100% of the Sepsis Screening Tool for their patients.  This review is to make sure that the initial decisions were appropriate to the final patient outcomes.					
MD Initials:	Time Zero:	Date:				